

CHAPTER 2.2.13.

(Background information on the proposed **Bluetongue** text below – The Terrestrial Code Commission discussed with the Scientific Commission Member Countries' comments on the revised chapter on bluetongue. The Terrestrial Code Commission examined comments received from Member Countries but made changes only when scientific justification accompanied those comments; this approach was endorsed by the Scientific Commission. The two Commissions were not aware of any new information to contradict the conclusions of the 2003 OIE Bluetongue Conference in Sicily regarding the infective period for bluetongue, and the Terrestrial Code Commission did not make any changes in this regard. The comment from Australia that the terminology relating to the *Culicoides* vector be modified to 'likely to support BTV transmission' was not adopted as the Terrestrial Code Commission considered that its proposed wording 'likely to be competent' adequately addressed the biology of the vector population and used accepted terminology. In response to comments from Australia and the EU, the period for which animals need to be vaccinated before movement was increased to 60 days to make it consistent with the accepted viraemic period. The comment from Peru regarding the point of departure in Article 2.2.13.6. was not adopted as the definition 'place of shipment' was considered to adequately cover the intent of the Article. After consulting with an expert, the Terrestrial Code Commission accepted changes recommended by the USA regarding the timing of serological testing in various articles relating to semen and embryo collection.)

BLUETONGUE

Article 2.2.13.1.

For the purposes of the *Terrestrial Code*, the *infective period* for bluetongue virus (BTV) shall be ~~400~~ 60 days.

The global BTV distribution ~~historically has been shown to be~~ is currently between latitudes of approximately ~~50~~⁴⁰°N and 35°S ~~but is known to be expanding in the northern hemisphere.~~

In the absence of clinical disease in a country or zone within this part of the world, its BTV status should be determined by an ongoing surveillance and monitoring programme (in accordance with ~~Chapter 1.3.6~~ Appendix 3.8.1) designed in accordance with the epidemiology of the disease, i.e. focusing on climatic and geographical factors, the biology and likely competence of *Culicoides* and/or serology of susceptible animals. The programme may need to be adapted to target parts of the country or zone at a higher risk due to historical, geographical and climatic factors, ruminant population data and *Culicoides* ecology, or proximity to enzootic or incursional zones as described in Appendix 3.8.1. ~~Random and targeted serological surveillance should provide at least a 95% level of confidence of detecting an annual seroconversion incidence of 2% in cattle (or other ruminant species if sufficient cattle are not available).~~

All countries or zones located outside this part of the world but adjacent to a country or zone not having free status should be subjected to similar surveillance. The surveillance programme should be carried out over a distance of at least 100 kilometres from the border with that country or zone, but a lesser distance could be acceptable if there are relevant ecological or geographical features likely to interrupt the transmission of BTV.

Standards for diagnostic tests and vaccines are described in the *Terrestrial Manual*.

Article 2.2.13.2.

BTV free country or zone

- 1) A country or a zone may be considered free from BTV when bluetongue is notifiable in the whole country and either:
 - a) the country or zone lies wholly north of ~~50~~⁴⁰°N or south of 35°S, and is not adjacent to a country or zone not having a free status; or
 - b) a surveillance and monitoring programme as described in ~~Chapter 1.3.6~~^{Appendix 3.8.1} ~~Article 2.2.13.1~~ has demonstrated no evidence of BTV in the country or zone during the past 2 years; ~~nor have any ruminants been vaccinated against bluetongue in the country or zone during the past 12 months;~~ or
 - c) a surveillance and monitoring programme has demonstrated no evidence of *Culicoides* likely to be competent BTV vectors in the country or zone.

~~For maintenance of the free status, the provisions of the last paragraph of Article 2.1.9.1. may need to be complied with on a continuous basis according to the geographical location of the country or zone.~~

- 2) A BTV free country or zone in which surveillance and monitoring has found no evidence that *Culicoides* likely to be competent BTV vectors are present will not lose its free status through the importation of vaccinated, seropositive or infective animals, or semen or embryos/ova from infected countries or zones.
- 3) A BTV free country or zone in which surveillance and monitoring has found evidence that *Culicoides* likely to be competent BTV vectors are present will not lose its free status through the importation of vaccinated or seropositive animals from infected countries or zones, provided:
 - a) the animals have been vaccinated in accordance with the *Terrestrial Manual* at least ~~30~~⁶⁰ days prior to dispatch with a vaccine which covers all serotypes whose presence in the source population has been demonstrated through a surveillance and monitoring programme as described in ~~Chapter 1.3.6~~^{Appendix 3.8.1}, and that the animals are identified in the accompanying certification as having been vaccinated; or
 - b) the animals are not vaccinated, and a surveillance and monitoring programme as described in ~~Chapter 1.3.6~~^{Appendix 3.8.1} has been in place in the source population for a period of 60 days immediately prior to dispatch, and no evidence of BTV transmission has been detected.
- 4) A BTV free country or zone adjacent to an infected country or zone should include a ~~surveillance~~ zone in which surveillance is conducted as described in ~~Appendix 3.8.1~~^{Article 2.1.9.1}. Animals within ~~this the surveillance~~ zone must be subjected to continuing surveillance. The boundaries of ~~the surveillance~~ this zone must be clearly defined, and must take account of geographical and epidemiological factors that are relevant to BTV transmission ~~infection~~.

Article 2.2.13.3.

BTV seasonally free zone

A BTV seasonally free zone is a part of an infected country or zone for which for part of a year, surveillance and monitoring demonstrate no evidence either of BTV transmission or of adult *Culicoides* likely to be competent BTV vectors.

For the application of Articles 2.2.13.7., 2.2.13.10. and 2.2.13.14., the seasonally free period is taken to commence the day following the last evidence of BTV transmission (as demonstrated by the surveillance and monitoring programme), or of the cessation of activity of adult *Culicoides* likely to be competent BTV vectors.

For the application of Articles 2.2.13.7., 2.2.13.10. and 2.2.13.14., the seasonally free period is taken to conclude either:

- 1) at least 28 days before the earliest date that historical data show bluetongue virus activity has recommenced; or
- 2) immediately if current climatic data or data from a surveillance and monitoring programme indicate an earlier resurgence of activity of adult *Culicoides* likely to be competent BTV vectors.

A BTV seasonally free zone in which surveillance and monitoring has found no evidence that *Culicoides* likely to be competent BTV vectors are present will not lose its free status through the importation of vaccinated, seropositive or infective animals, or semen or embryos/ova from infected countries or zones.

Article 2.2.13.4.

BTV infected country or zone

A BTV infected country or zone is a clearly defined area where evidence of BTV has been reported during the past 2 years.

Article 2.2.13.5.

Veterinary Administrations of countries shall consider whether there is a risk with regard to BTV infection in accepting importation or transit through their territory, from other countries, of the following *commodities*:

- 1) ruminants and other BTV susceptible herbivores;
- 2) semen of these species;
- 3) embryos/ova of these species;
- 4) *pathological material* and biological products (from these species) (see Chapter 1.4.6. and Section 1.5.).

Other *commodities* should be considered as not having the potential to spread BTV when they are the subject of *international trade*.

Article 2.2.13.6.

When importing from BTV free countries or zones, *Veterinary Administrations* should require:

for ruminants and other BTV susceptible herbivores

the presentation of an *international veterinary certificate* attesting that:

- 1) the animals were kept in a BTV free country or zone since birth or for at least ~~60~~400 days prior to shipment; or
- 2) the animals were kept in a BTV free country or zone for at least 28 days, then were subjected, with negative results, to a serological test to detect antibody to the BTV group according to the *Terrestrial Manual*, ~~such as the BT competition ELISA or the BT AGID test~~, and remained in the BTV free country or zone until shipment; or
- 3) the animals were kept in a BTV free country or zone for at least 7 days, then were subjected, with negative results, to an agent identification test according to the *Terrestrial Manual* ~~a BTV isolation test or polymerase chain reaction test on a blood sample~~, and remained in the BTV free country or zone until shipment; or
- 4) the animals:
 - a) were kept in a BTV free country or zone for at least 7 days;
 - b) were vaccinated in accordance with the *Terrestrial Manual* ~~30~~ 60 days before introduction into the free country or zone against all serotypes whose presence in the source population has been demonstrated through a surveillance and monitoring programme as described in Appendix 3.8.1.;
 - c) were identified as having been vaccinated and
 - d) remained in the BTV free country or zone until shipment;
- 5)
 - 4) if the animals were exported from a free zone, either:
 - a) did not transit through an infected zone during transportation to the *place of shipment*; or
 - b) were protected from attack from *Culicoides* likely to be competent BTV vectors at all times when transiting through an infected zone; or
 - c) had been vaccinated in accordance with point 4) above.

Article 2.2.13.7.

When importing from BTV seasonally free zones, *Veterinary Administrations* should require:

for ruminants and other BTV susceptible herbivores

the presentation of an *international veterinary certificate* attesting that the animals:

- 1) were kept during the seasonally free period in a BTV seasonally free zone for at least ~~60~~400 days prior to shipment; or
- 2) were kept during the BTV seasonally free period in a BTV seasonally free zone for at least 28 days prior to shipment, and were subjected during the residence period in the zone to a serological test to detect antibody to the BTV group, according to the *Terrestrial Manual* ~~such as~~

~~the BT competition ELISA or the BT AGID test~~, with negative results on two occasions, with an interval of not less than 7 days between each test, the first test being carried out at least 21 days after the commencement of the residence period; or

- 3) were kept during the BTV seasonally free period in a BTV seasonally free zone for at least 14 days prior to shipment, and were subjected during the residence period in the zone to an agent identification test according to the Terrestrial Manual ~~to a BTV isolation test or polymerase chain reaction test~~, with negative results, on blood samples taken on two occasions, with an interval of not less than 7 days between each test, the first test being carried out at least 7 days after the commencement of the residence period; or
- 4) were kept during the seasonally free period in a BTV seasonally free zone, and were vaccinated in accordance with the Terrestrial Manual 30 60 days before introduction into the free country or zone against all serotypes whose presence in the source population has been demonstrated through a surveillance and monitoring programme as described in Appendix 3.8.1, were identified as having been vaccinated and remained in the BTV free country or zone until shipment;

AND

~~5)~~4) if the animals were exported from a free zone, either:

- a) did not transit through an infected zone during transportation to the *place of shipment*, or
- b) were protected from attack from *Culicoides likely to be competent BTV vectors* at all times when transiting through an infected zone, or
- c) were vaccinated in accordance with point 4) above.

Article 2.2.13.8.

When importing from BTV infected countries or zones, *Veterinary Administrations* should require:

for ruminants and other BTV susceptible herbivores

the presentation of an *international veterinary certificate* attesting that the animals:

- 1) were protected from attack from *Culicoides likely to be competent BTV vectors* for at least ~~60~~40 days prior to shipment; or
- 2) were protected from attack from *Culicoides likely to be competent BTV vectors* for at least 28 days prior to shipment, and were subjected during that period to a serological test according to the Terrestrial Manual to detect antibody to the BTV group, ~~such as the BT competition ELISA or the BT AGID test~~, with negative results on two occasions, with an interval of not less than 7 days between each test, the first test being carried out at least 21 days after introduction into the *quarantine station*; or
- 3) were protected from attack from *Culicoides likely to be competent BTV vectors* for at least 14 days prior to shipment, and were subjected during that period to an agent identification test according to the Terrestrial Manual ~~a BTV isolation test or polymerase chain reaction test~~, with negative results, on blood samples taken on two occasions, with an interval of not less than 7 days between each test, the first test being carried out at least 7 days after introduction into the *quarantine station*; or

- 4) were vaccinated in accordance with the *Terrestrial Manual* at least ~~30~~ 60 days before shipment, against all serotypes whose presence in the source population has been demonstrated through a surveillance and monitoring programme as described in Appendix 3.8.1, and were identified in the accompanying certification as having been vaccinated; or
- 5) are not vaccinated, a surveillance and monitoring programme as described in Appendix 3.8.1. has been in place in the source population for a period of 60 days immediately prior to shipment, and no evidence of BTV transmission has been detected;

AND

- 6) were protected from attack from Culicoides likely to be competent BTV vectors during transportation to the *place of shipment*; or
- 7) were vaccinated ~~30~~ 60 days before shipment or had antibodies against all serotypes whose presence in the zones of transit has been demonstrated through a surveillance and monitoring programme as described in Appendix 3.8.1.

Article 2.2.13.9.

When importing from BTV free countries or zones, *Veterinary Administrations* should require:

for semen of ruminants and other BTV susceptible herbivores

the presentation of an *international veterinary certificate* attesting that:

- 1) the donor animals:
 - a) were kept in a BTV free country or zone for at least ~~60~~400 days before commencement of, and during, collection of the semen; or
 - b) were subjected to a serological test according to the *Terrestrial Manual* to detect antibody to the BTV group, ~~such as the BT competition ELISA or the BT AGID test,~~ between ~~28~~ 21 and 60 days after the last collection for this consignment, with negative results; or
 - c) were subjected to an agent identification test according to the *Terrestrial Manual* ~~a virus isolation test or polymerase chain reaction (PCR) test~~ on blood samples collected at commencement and conclusion of, and at least every 7 days (virus isolation test) or at least every 28 days (PCR test) during, semen collection for this consignment, with negative results;
- 2) the semen was collected, processed and stored in conformity with the provisions of either Appendix 3.2.1. or Appendix 3.2.2.

Article 2.2.13.10.

When importing from BTV seasonally free zones, *Veterinary Administrations* should require:

for semen of ruminants and other BTV susceptible herbivores

the presentation of an *international veterinary certificate* attesting that:

- 1) the donor animals:
 - a) were kept during the BTV seasonally free period in a seasonally free zone for at least

~~60~~⁴⁰ days before commencement of, and during, collection of the semen; or

- b) were subjected to a serological test according to the *Terrestrial Manual* to detect antibody to the BTV group ~~such as the BT competition ELISA or the BT AGID test~~, with negative results, at least every 60 days throughout the collection period and between ~~28~~ 21 and 60 days after the final collection for this consignment; or
 - c) were subjected to an agent identification test according to the *Terrestrial Manual* ~~a virus isolation test or polymerase chain reaction (PCR) test~~ on blood samples collected at commencement and conclusion of, and at least every 7 days (virus isolation test) or at least every 28 days (PCR test) during, semen collection for this consignment, with negative results;
- 2) the semen was collected, processed and stored in conformity with the provisions of either Appendix 3.2.1. or Appendix 3.2.2.

Article 2.2.13.11.

When importing from BTV infected countries or zones, *Veterinary Administrations* should require:

for semen of ruminants and other BTV susceptible herbivores

the presentation of an *international veterinary certificate* attesting that:

- 1) the donor animals:
 - a) were protected from attack from *Culicoides* likely to be competent BTV vectors for at least ~~60~~⁴⁰ days before commencement of, and during, collection of the semen; or
 - b) were subjected to a serological test according to the *Terrestrial Manual* to detect antibody to the BTV group ~~such as the BT competition ELISA or the BT AGID test~~, with negative results, at least every 60 days throughout the collection period and between ~~28~~ 21 and 60 days after the final collection for this consignment; or
 - c) were subjected to an agent identification test according to the *Terrestrial Manual* ~~a virus isolation test or polymerase chain reaction (PCR) test~~ on blood samples collected at commencement and conclusion of, and at least every 7 days (virus isolation test) or at least every 28 days (PCR test) during, semen collection for this consignment, with negative results;
- 2) the semen was collected, processed and stored in conformity with the provisions of either Appendix 3.2.1. or Appendix 3.2.2.

Article 2.2.13.12.

Regardless of the bluetongue status of the *exporting country*, *Veterinary Administrations* of *importing countries* should require:

for *in vivo* derived bovine embryos/oocytes

the presentation of an *international veterinary certificate* attesting that the embryos/oocytes were collected, processed and stored in conformity with the provisions of Appendix 3.3.1. or Appendix 3.3.3., as relevant.

Article 2.2.13.13.

When importing from BTV free countries or zones, *Veterinary Administrations* should require:
for *in vivo* derived embryos of ruminants (other than bovines) and other BTV susceptible herbivores
the presentation of an *international veterinary certificate* attesting that:

- 1) the donor females:
 - a) were kept in a BTV free country or zone for at least the ~~60~~400 days prior to, and at the time of, collection of the embryos; or
 - b) were subjected to a serological test according to the *Terrestrial Manual* to detect antibody to the BTV group, ~~such as the BT competition ELISA or the BT AGID test~~, between ~~28~~ 21 and 60 days after collection, with negative results; or
 - c) were subjected to an agent identification test according to the *Terrestrial Manual* ~~a BTV isolation test or polymerase chain reaction test~~ on a blood sample taken on the day of collection, with negative results;
- 2) the embryos were collected, processed and stored in conformity with the provisions of Appendix 3.3.1.

Article 2.2.13.14.

When importing from BTV seasonally free zones, *Veterinary Administrations* should require:
for *in vivo* derived embryos/oocytes of ruminants (other than bovines) and other BTV susceptible herbivores and for *in vitro* produced bovine embryos
the presentation of an *international veterinary certificate* attesting that:

- 1) the donor females:
 - a) were kept during the seasonally free period in a seasonally free zone for at least ~~60~~400 days before commencement of, and during, collection of the embryos/oocytes; or
 - b) were subjected to a serological test according to the *Terrestrial Manual* to detect antibody to the BTV group, ~~such as the BT competition ELISA or the BT AGID test~~, between ~~28~~ 21 and 60 days after collection, with negative results; or
 - c) were subjected to an agent identification test according to the *Terrestrial Manual* ~~a BTV isolation test or polymerase chain reaction test~~ on a blood sample taken on the day of collection, with negative results;
- 2) the embryos/oocytes were collected, processed and stored in conformity with the provisions of Appendix 3.3.1.

Article 2.2.13.15.

When importing from BTV infected countries or zones, *Veterinary Administrations* should require:
for *in vivo* derived embryos/oocytes of ruminants (other than bovines) and other BTV susceptible herbivores and for *in vitro* produced bovine embryos
the presentation of an *international veterinary certificate* attesting that:

- 1) the donor females:
 - a) were protected from attack from *Culicoides* likely to be competent BTV vectors for at least

- ~~60400~~ days before commencement of, and during, collection of the embryos/oocytes; or
- b) were subjected to a serological test according to the *Terrestrial Manual* to detect antibody to the BTV group, ~~such as the BT competition ELISA or the BT AGID test~~, between ~~28~~ 21 and 60 days after collection, with negative results; or
 - c) were subjected to an agent identification test according to the *Terrestrial Manual* ~~a BTV isolation test or polymerase chain reaction test~~ on a blood sample taken on the day of collection, with negative results;
- 2) the embryos/oocytes were collected, processed and stored in conformity with the provisions of Appendix 3.3.1.

Article 2.2.13.16.

Protecting animals from Culicoides attack

When transporting animals through BTV infected countries or zones, Veterinary Administrations should require strategies to protect animals from attack from *Culicoides* likely to be competent BTV vectors during transport, taking into account the local ecology of the vector.

Strategies to protect animals from attack from *Culicoides* likely to be competent BTV vectors during transport through an infected country or zone should take into account the local ecology of the vector.

Potential risk management strategies include:

- 1) treating animals with chemical repellents prior to and during transportation;
- 2) loading, transporting and unloading animals at times of low vector activity i.e. bright sunshine, low temperature;
- 3) ensuring vehicles do not stop en route during dawn or dusk, or overnight, unless the animals are held behind insect proof netting;
- 4) darkening the interior of the vehicle, for example by covering the roof and/or sides of vehicles with shade cloth;
- 5) monitoring for vectors at common stopping and offloading points to gain information on seasonal variations;
- 6) using historical, ongoing and/or BTV modeling information to identify low risk ports and transport routes.